

1 Convergent evolution of increased urine concentrating ability in desert mammals

2

3 **Authors:** Joana L. Rocha^{1,2}, José C. Brito^{1,2}, Rasmus Nielsen^{3,4} and Raquel Godinho^{1,2,5}

4

5 ¹CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Campus de Vairão, 4485-
6 661 Vairão, Portugal.

7 ²Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, 4169-007 Porto, Portugal.

8 ³Department of Integrative Biology and Department of Statistics, University of California Berkeley, Berkeley, CA 94820, USA.

9 ⁴Globe Institute, University of Copenhagen, DK-1165 Copenhagen, Denmark.

10 ⁵Department of Zoology, University of Johannesburg, PO Box 534, Auckland Park 2006, South Africa.

11

12 **Author for correspondence:** Joana L. Rocha, email: joana.laranjeira.rocha@gmail.com; Raquel
13 Godinho, email: rgodinho@cibio.up.pt;

14

15 Abstract

16

17 One of the most celebrated textbook examples of physiological adaptations to desert
18 environments is the unique ability that desert mammals have to produce hyperosmotic urine.
19 Commonly perceived as an adaptation mainly observed in small rodents, the extent to which urine
20 concentrating ability has independently evolved in distinct lineages, including medium-sized and
21 large desert mammals, has not previously been assessed using modern phylogenetic approaches.
22 Here, we explicitly test the general hypothesis that desert-dwelling mammals have evolved
23 increased ability to concentrate urine compared to non-desert species, controlling for body mass
24 and other covariates. Phylogenetic generalized least-squares models show that the mean aridity
25 index of a species distribution range largely predicts its urine concentrating ability, even when
26 accounting for body mass differences and phylogenetic correlations. In contrast, we find much
27 weaker correlations between mass-adjusted basal metabolic rate and environmental variables.

28

29 **Subject areas:** ecology, evolution, comparative physiology, desert biology

30

31 **Keywords:** phylogenetic generalized least squares, adaptation, hyperosmotic urine, deserts,
32 aridity index

33

34

35

36

37

38 1. Background

39

40 Deserts are defined as regions with aridity index, a ratio of annual precipitation and
41 potential evapo-transpiration, below 0.20 [1]. Desert species independently evolved striking
42 adaptations to cope with the water scarcity and extreme climatic and physical conditions that
43 characterize such habitats [2–7]. For the last fifty years, this has captured the attention of many
44 eco-physiologists who turned to desert biology to study the physiological mechanisms that help
45 maintaining body temperature and retain water. Remarkably, most of the classic works on
46 mammalian desert physiology, pioneered by Schmidt-Nielsen et al [8–11], have survived the test
47 of time, as researchers have applied new perspectives and tools to test specific hypothesis
48 regarding the evolution of adaptive traits [12,13].

49 Among the classical findings of physiological adaptations that minimize water loss, is the
50 ability of some desert mammals to produce highly concentrated urine [14]. This phenotype is
51 mostly associated with desert ‘evaders’, such as the Australian Hopping mouse, which has the
52 record for highest hyperosmotic urine (above 9000 mOsm/kg) [3,15]. Evaders, part of a
53 classification system proposed by Willmer *et al.* (2000), are small body-sized animals that are able
54 to evade extreme conditions through behavior [5]. By contrast, large-sized mammals unable to
55 shelter from extreme climates that are forced to withstand heat, are called ‘endurers’, and
56 medium-sized mammals unable to evade nor withstand extremes as efficiently as evaders and
57 endurers are called ‘evaporators’ [5]. Because mammalian urine concentrating ability (mOsm/Kg)
58 is negatively correlated with body mass [16], desert evaders stand out in this capacity [15].
59 However, when correcting for differences in body mass, the capacity to highly concentrate urine
60 seems to have independently evolved in deserts. For example, evaporators such as the fennec fox
61 can concentrate urine up to 4022 mOsm/Kg, which is much more concentrated than in similar-
62 sized non-desert counterparts and, when mass-adjusted, nearly as impressive as the 5500
63 mOsm/Kg of a lesser Egyptian gerbil [5,17,18].

64 Classical comparative physiology studies have provided evidence for convergent adaptive
65 evolution of ecologically relevant phenotypes in multiple desert mammals. A textbook example
66 includes basal metabolic rate (BMR), which is positively correlated with zoogeographical zones,
67 even when accounting for phylogeny and body mass [19,20]. Desert mammals evolved lower
68 production of metabolic heat (low BMR) to maintain body temperatures and avoid water loss by
69 evaporative cooling [19–23]. Surprisingly, and to the best of our knowledge, no study to date has
70 attempted to identify continuous climatic or environmental variables that help explain variation in
71 urine concentrating ability, one of the most celebrated examples of mammalian desert adaptation.
72 Moreover, and in contrast to BMR, previous studies concerning urine concentrating ability have

73 ignored phylogenetic correlations when analyzing associations between this trait and the
74 environment [16,24].

75 Here, we explicitly test the hypothesis that the aridity index of a species range predicts
76 maximum urine concentrating ability, even when accounting for body mass and phylogenetic
77 relationship, and provide statistical evidence that the ability to avoid water loss by producing
78 hyperosmotic urine has evolved independently in multiple phylogenetic lineages of desert
79 mammals. Additionally, we use similar models to re-analyze mass adjusted BMR for species in
80 which urine osmolality was obtained, and test aridity index and temperature variables as
81 predictors for metabolism in deserts.

82

83 2. Methods

84

85 Phenotypes were obtained by extensive literature search and by taking advantage of large
86 datasets from previously published revision studies [16,19,24,25]. A total of 108 mammalian
87 species' maximum urine concentrating abilities came from two previous studies by Beuchat
88 [16,24], which recorded the specific condition (factor defined as C) under which urine osmolality
89 (hereafter mOsm) was measured, including: dehydrated (D), given salt (S) or protein (P) loading,
90 combinations of these conditions (DP, DS, SP, DSP), or treatment not specified (X), as well as the
91 method used to measure urine mOsm (see 15,23 for more information). We also recorded the
92 highest urine osmolality values for 13 additional species from other studies [18,26–30], and also
93 noted the experimental conditions (C) the study-cases were subjected to, the method (M) used to
94 measure urine mOsm, and the study-species diet (Di), following the same notation as Beuchat (see
95 15,23 for more information). For species with more than one record of urine mOsm, due to
96 different treatments/studies, we opted to keep values taken from captive study-subjects under
97 controlled/experimentally induced dehydration, as opposed to measurements of unknown
98 condition taken from the field. Ultimately, only one individual was used per species.

99 As mOsm is a plastic trait, the experimental conditions under which this parameter is
100 measured can greatly affect estimates. In particular, the degree of hydration of the study animals
101 naturally affects urine osmolality. As such, we curated two datasets: 1) a larger dataset including
102 all 121 reported measurements of mOsm from Beuchat [16,24] and our search [18,26–30]; 2) a
103 subset of the larger dataset only containing 87 observations in which the study-subject, in
104 captivity, was dehydrated (D) and not given food or any other treatment. However, we note that
105 there is still some residual variation in the number of days under water deprivation for an animal
106 to be considered fully dehydrated and variation in the method used for measuring mOsm. Also,
107 for the 121 species dataset, 24 records came from studies that did not report the procedures used

108 to measure mOsm. A full detail of the conditions and studies can be found in electronic
109 Supplementary Material TableS1. Downstream analyses were performed on both data sets.

110 The distribution polygons of each species for which physiological data were available were
111 downloaded from the IUCN database [31,32]. We also downloaded the following environmental
112 variables at about 1km spatial resolution: annual aridity index (AI) from CGIAR-CSI Global Aridity
113 database [33], and annual mean temperature (BIO1), maximum temperature of the warmest
114 month (BIO5), maximum temperature of the coldest month (BIO6), and temperature annual range
115 (BIO7) from the WorldClim database [34]. We then used a Geographical Information System to
116 calculate mean values for each environmental variable within the distribution polygon of each
117 species. For species with global distributions ranges, encompassing humid to arid environments,
118 and for which phenotypes under review were sampled from a specific region, we partitioned the
119 original distribution polygon accordingly and calculated mean values for the sampled zone. To
120 avoid the effects of collinearity we tested for correlations between pairs of variables using a
121 Spearman correlation test before testing them together as predictors for mOsm and BMR in
122 downstream analyses. We used a previously published mammalian tree estimated using both
123 nuclear and mitochondrial genes [35] and pruned the tree to only retain the species included in
124 or study. For visualization of trait evolution on the tree, we projected states for maximum mOsm
125 and bioclimatic variables, estimated using maximum likelihood, onto the internal edges and nodes
126 of the tree using a color gradient. The estimation was done using the *fastAnc* option in the
127 'contMap' function implemented in 'phytools' [36].

128 Phylogenetic generalized least squares (PGLS) as implemented in 'phytools' were
129 performed using the 'gls' function with the model of Pagel (1999) [37] to test different linear
130 models with increasing complexity. Briefly, the model of Pagel adjusts the off-diagonal elements
131 of variance-covariance matrix in a Brownian evolution model with a multiplicative factor, λ , such
132 that when $\lambda = 0$ a star phylogeny is obtained representing no phylogenetic signal, while when $\lambda =$
133 1 a standard Brownian on the reference tree is obtained. As residuals of maximum urine osmolality
134 are not normally distributed (**Figure S1**), we performed PGLS analyses on $\log_{10}(\text{maximum mOsm})$
135 using $\log_{10}(\text{body mass})$ as covariate. Mean annual aridity index (hereafter mean AI) and
136 uncorrelated mean temperature variables were used in PGLS models as potential predictors of
137 urine concentrating ability. Condition (C), method (M) and diet (Di) were also included as
138 covariates. For the dataset including only dehydrated individuals, the same PGLS models were
139 tested with AI converted to a categorical variable using the following binning defined in [33]:
140 humid for $\text{AI} > 0.65$, dry sub-humid for $0.5 < \text{AI} < 0.65$, semi-arid for $0.2 < \text{AI} < 0.5$, and arid for AI
141 < 0.2 . To test for phylogenetic signal for specific traits, we calculated Pagel's λ on urine
142 concentrating ability, body mass and AI using the function 'phylosig' as implemented in 'phytools'.

143 3. Results

144

145 We analyzed a dataset with information on maximum urine concentrating abilities, mean
146 annual aridity index and WorldClim temperatures for a total of 121 species of mammals, with
147 available DNA sequence information, spanning 6 g to 479 kg of body mass. Using the classification
148 of [33] the dataset contained 32, 35, 15, and 39 species with distribution ranges classified as arid,
149 semi-arid, dry sub-humid, and humid, respectively. Ancestral state reconstruction across our
150 mammalian dataset's phylogeny shows that the ability to produce hyperosmotic urine has evolved
151 multiple times in species with distribution ranges characterized by low mean AI (**Figure 1**).

152 When testing for phylogenetic signal using Pagel's λ we found highly significant
153 phylogenetic signal for most variables, validating the relevance of adjusting the linear models for
154 phylogenetic covariance (**Table 1**). The mean AI correlates with all temperature variables, except
155 mean annual temperature range (**Figure S3**). To avoid collinearity, all WorldClim temperatures but
156 the latter were excluded from PGLS analysis with mean AI (**Table S2**).

157 All PGLS models tested in this study point to mean annual aridity index as a highly
158 significant and important predictor of mammalian maximum urine concentration ability (**Table 2**,
159 **TableS2**). The lower the mean AI, the more efficient mammals are able to concentrate urine
160 (**Figure 2, FigureS2**), even when accounting for the differences in body mass, diet, experimental
161 condition, methodology and phylogeny (**Table 2**). Overall, species inhabiting arid and semi-arid
162 environments can maximize urine concentration well above the levels of dry sub-humid and humid
163 counterparts (**Figure 2**). In congruence with past studies, our results also show that log-
164 transformed body mass is also negatively correlated with log-transformed maximum urine
165 osmolality [16], though not to the same extent as mean aridity index (**Table 2, Figure S2**). Diet is
166 also a significant predictor of maximum urine osmolality in fructivorous mammals, which
167 significantly concentrate more urine than strictly carnivore species (**Table 2**). Overall, no strong
168 correlation was found between urine concentrating ability and temperature variables (**Figure S4**,
169 **TableS2**).

170 When the experimental condition was used as co-variate in the PGLS analysis, its role in
171 predicting maximum urine osmolality was not significant (**Table 2, Table S2**). Nonetheless, we also
172 tested similar PGLS models on a subset of the data that only included individuals fulfilling the
173 dehydrated criterion (see Methods section), resulting in a smaller dataset including 87 species (18
174 arid, 26 semi-arid, 13 dry sub-humid, and 30 humid). The results from this analysis are similar to
175 those of the larger dataset conditions (**Table 2**). When using AI as a discrete variable we found
176 that urine concentrating ability is significantly higher in species from arid environments than in
177 species from semi-arid, dry sub-humid and humid regions (**Table S3**). In all analyses, the aridity

178 index of the species distribution is a stronger predictor of maximum urine osmolality than body
179 mass and diet. Notably, experimental condition and methodology are non-significant.

180 We also obtained mass-adjusted basal metabolic rates (BMR) for 84 of the 121 species
181 from our dataset. BMR is weakly correlated with bioclimatic variables (**Figure S5**), and differences
182 in this parameter are better explained by log₁₀ body mass (**TableS4**).

183

184 4. Discussion

185

186 The extent to which different mammalian species from desert environments have
187 significantly higher urine concentrating abilities, a phenotype to better retain water in extreme
188 conditions of water-deprivation and extreme temperature ranges, has not previously been subject
189 to rigorous statistical analyses that account for phylogeny. Here we show that increased ability to
190 concentrate urine, measured as maximum urine osmolality, has convergently evolved in many
191 desert mammals (**Figure 1**).

192 PGLS analysis on a multi-species dataset revealed that desert-dwelling species effectively
193 concentrate more urine than non-desert species, even when accounting for both body mass and
194 ancestry, and furthermore, that differences in mammalian maximum urine concentrating ability
195 can be predicted, to some degree, by the aridity of species' range. Our measurement of
196 environmental aridity is inherently imprecise as it does not account for spatial and temporal
197 changes in the environmental conditions nor in the species distributions, and it implicitly assumes
198 a uniform density distribution across the species range [33,34]. Still, we find it more rigorous than
199 the often-used binary approach of classifying species as 'mesic' or 'xeric' based on eye-assessment
200 of species distribution range or habitat descriptions (e.g. [19,24]). Despite these challenges, and
201 while acknowledging additional noise added by varying experimental conditions and
202 methodologies, we find a strongly significant correlation between maximum urine osmolality and
203 mean annual aridity index. This observation is similar to what has been previously reported in
204 other desert adaptive phenotypes such as basal metabolic rate [19,20]. However, we find a much
205 weaker correlation for BMR and environmental variables when analyzing 84 of the same species
206 as in the mOsm analysis, suggesting that, in contrast to our results for mOsm, differences in BMR
207 among species are mostly explained by body mass differences and are only weakly predicted by
208 environmental variables [38].

209 A contentious debate in the fields of ecological and evolutionary physiology is whether
210 phenotypic differences between species and populations inhabiting contrasting environments are
211 due to genetic (evolutionary) adaptations or due to plastic responses, and difference in the ability
212 to concentrate urine is no exception. Although all mammals, when water-deprived, are able to

213 increase urine concentration in relation to their optimal hydrated states [14,39], our results for a
214 dataset in which species were all captive and water-deprived, provide strong evidence that
215 increased maximal urine concentration is an adaptation to high aridity and has evolved multiple
216 times in different phylogenetic lineages. The hypothesis that higher maximum urine concentrating
217 ability in mammals is an adaptive trait is also supported by an observed increase in renal medullary
218 thickness in desert mammals [3,16].

219 Our PGLS analyses also suggest that, among the many physical and climatic challenges
220 faced by desert species, aridity, more than temperature, has been one of the main selective
221 pressures increasing maximum urine concentrating ability, and driving its repeated evolution in
222 different desert mammalian lineages. As population and comparative genomics studies in desert
223 environments continue to search for the genetic basis of desert adaptation (e.g. [40–43]),
224 maximum urine concentrating ability might be one of the primary physiological traits that lends
225 itself to more detailed genetic analyses.

226

227

228 **Author's contributions:** J.L.R contributed to the conception and design of the study, data
229 collection, analysis, and drafted the manuscript. R.G, J.C.B, R.N contributed to the study's
230 conception and design, interpretation of data, editing and critical revision. All authors approve
231 publication and agree to be accountable for all aspects of the work.

232

233 **Acknowledgments:** We thank Pedro Tarroso and Antigoni Kaliontzopoulou for discussions that
234 helped conceiving this study.

235

236 **Funding:** J.L.R., J.C.B and R.G. were supported by the Portuguese Foundation for Science and
237 Technology, FCT (SFRH/BD/116397/2016, CEECINST/00014/2018, and contract under DL57/2016,
238 respectively). This work was partially supported by FCT project PTDC/BIA-EVL/31902/2017.

239

240 **Competing interests:** We declare we have no competing interests

241

242 **Ethics:** No ethical approval was required as this work only uses publicly available data.

243

244

245

246

247


248 REFERENCES

- 249 1. Ward D. 2016 *The biology of deserts*. 2nd edn. Oxford University Press.
- 250 2. Nagy KA. 2004 Water economy of free-living desert animals. *Int. Congr. Ser.* **1275**, 291–
- 251 297. (doi:10.1016/j.ics.2004.08.054)
- 252 3. Schwimmer H, Haim A. 2009 Physiological adaptations of small mammals to desert
- 253 ecosystems. *Integr. Zool.* **4**, 357–366. (doi:10.1111/j.1749-4877.2009.00176.x)
- 254 4. Fuller A, Hetem RS, Maloney SK, Mitchell D. 2014 Adaptation to heat and water shortage
- 255 in large, arid-zone mammals. *Physiology* **29**, 159–167. (doi:10.1152/physiol.00049.2013)
- 256 5. Willmer P, Stone G, Johnston IA. 2005 *Environmental physiology of animals*. 2nd edn.
- 257 Blackwell Pub.
- 258 6. Pannabecker TL. 2013 Comparative physiology and architecture associated with the
- 259 mammalian urine concentrating mechanism: Role of inner medullary water and urea
- 260 transport pathways in the rodent medulla. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.*
- 261 **304**, 488–503. (doi:10.1152/ajpregu.00456.2012)
- 262 7. Rymer TL, Pillay N, Schradin C. 2016 Resilience to droughts in mammals: A conceptual
- 263 framework for estimating vulnerability of a single species. *Q. Rev. Biol.* **91**, 133–176.
- 264 (doi:10.1086/686810)
- 265 8. SCHMIDT-NIELSEN B, O'DELL R. 1961 Structure and concentrating mechanism in the
- 266 mammalian kidney. *Am. J. Physiol.* **200**, 1119–1124.
- 267 (doi:10.1152/ajplegacy.1961.200.6.1119)
- 268 9. Schmidt-nielsen K. 1965 Desert animals. Physiological problems of heat and water. *Desert*
- 269 *Anim. Physiol. Probl. heat water*.
- 270 10. Schmidt-nielsen B, Schmidt-Nielsen K, Brokaw A, Schneiderman H. 1948 Water
- 271 conservation in desert rodents. *J. Cell. Comp. Physiol.* **32**, 331–360.
- 272 (doi:10.1002/jcp.1030320306)
- 273 11. Schmidt-Nielsen K, Haines HB, Hackel DB. 1964 Diabetes mellitus in the sand rat induced
- 274 by standard laboratory diets. *Science (80-.)*. **143**, 689–690.
- 275 (doi:10.1126/science.143.3607.689)
- 276 12. Tracy RL, Walsberg GE. 2002 Kangaroo rats revisited: Re-evaluating a classic case of desert
- 277 survival. *Oecologia* **133**, 449–457. (doi:10.1007/s00442-002-1059-5)
- 278 13. Walsberg GE. 2000 Small mammals in hot deserts: Some generalizations revisited.
- 279 *Bioscience*. **50**, 109–120. (doi:10.1641/0006-3568(2000)050[0109:SMIHDS]2.3.CO;2)
- 280 14. Sands JM, Layton HE. 2014 Advances in Understanding the Urine-Concentrating
- 281 Mechanism. *Annu. Rev. Physiol.* **76**, 387–409. (doi:10.1146/annurev-physiol-021113-
- 282 170350)
- 283 15. Donald J, Pannabecker TL. 2015 Osmoregulation in Desert-Adapted Mammals. In *Sodium*
- 284 *and Water Homeostasis*, pp. 191–211. Springer New York. (doi:10.1007/978-1-4939-3213-
- 285 9_10)
- 286 16. Beuchat CA. 1990 Body size, medullary thickness, and urine concentrating ability in
- 287 mammals. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **258**.
- 288 (doi:10.1152/ajpregu.1990.258.2.r298)
- 289 17. Noll-Banholzer U. 1979 Water balance and kidney structure in the fennec. *Comp.*
- 290 *Biochem. Physiol. -- Part A Physiol.* **62**, 593–597. (doi:10.1016/0300-9629(79)90109-9)

- 291 18. Kronfeld N, Shkolnik A. 1996 Adaptation to Life in the Desert in the Brown Hare (*Lepus*
292 *capensis*). *J. Mammal.* **77**, 171–178. (doi:10.2307/1382718)
- 293 19. Lovegrove BG. 2000 The zoogeography of mammalian basal metabolic rate. *Am. Nat.* **156**,
294 201–219. (doi:10.1086/303383)
- 295 20. Fristoe TS, Burger JR, Balk MA, Khaliq I, Hof C, Brown JH. 2015 Metabolic heat production
296 and thermal conductance are mass-independent adaptations to thermal environment in
297 birds and mammals. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 15934–15939.
298 (doi:10.1073/pnas.1521662112)
- 299 21. Lovegrove BG. 2003 The influence of climate on the basal metabolic rate of small
300 mammals: A slow-fast metabolic continuum. *J. Comp. Physiol. B Biochem. Syst. Environ.*
301 *Physiol.* **173**, 87–112. (doi:10.1007/s00360-002-0309-5)
- 302 22. Clarke A, Rothery P, Isaac NJB. 2010 Scaling of basal metabolic rate with body mass and
303 temperature in mammals. *J. Anim. Ecol.* **79**, 610–619. (doi:10.1111/j.1365-
304 2656.2010.01672.x)
- 305 23. Elgar MA, Harvey PH. 1987 Basal Metabolic Rates in Mammals: Allometry, Phylogeny and
306 Ecology. *Funct. Ecol.* **1**, 25. (doi:10.2307/2389354)
- 307 24. Beuchat CA. 1996 Structure and concentrating ability of the mammalian kidney:
308 Correlations with habitat. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **271**.
309 (doi:10.1152/ajpregu.1996.271.1.r157)
- 310 25. White CR, Seymour RS. 2003 Mammalian basal metabolic rate is proportional to body
311 mass^{2/3}. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 4046–4049. (doi:10.1073/pnas.0436428100)
- 312 26. Maloiy G, Macfarlane W, Shkolnik A. 1979 Mammalian herbivores.
- 313 27. Diaz GB, Ojeda RA. 1999 Kidney structure and allometry of Argentine desert rodents. *J.*
314 *Arid Environ.* **41**, 453–461. (doi:10.1006/jare.1998.0472)
- 315 28. Nagy KA, Shoemaker VH, Costa WR. 1976 Water, Electrolyte, and Nitrogen Budgets of
316 Jackrabbits (*Lepus californicus*) in the Mojave Desert. *Physiol. Zool.* **49**, 351–363.
317 (doi:10.1086/physzool.49.3.30155693)
- 318 29. Ostrowski S, Williams JB, Mésochina P, Sauerwein H. 2006 Physiological acclimation of a
319 desert antelope, Arabian oryx (*Oryx leucoryx*), to long-term food and water restriction. *J.*
320 *Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **176**, 191–201. (doi:10.1007/s00360-005-
321 0040-0)
- 322 30. Downs CT, Perrin MR. 1991 Urinary concentrating ability of four Gerbillurus species of
323 southern African arid regions. *J. Arid Environ.* **20**, 71–81. (doi:10.1016/s0140-
324 1963(18)30776-6)
- 325 31. IUCN. 2020 International Union for Conservation of Nature. The IUCN Red List of
326 Threatened Species. Version 2020-1.
- 327 32. IUCN. 2019 International Union for Conservation of Nature. The IUCN Red List of
328 Threatened Species. Version 2019-1.
- 329 33. Trabucco A, Zomer R. 2009 Global Aridity Index (Global-Aridity) and Global Potential
330 Evapo-Transpiration (Global-PET) Geospatial Database. *CGIAR Consort. Spat. Inf.*
331 (doi:10.1017/CBO9781107415324.004)
- 332 34. Fick SE, Hijmans RJ. 2017 WorldClim 2: new 1-km spatial resolution climate surfaces for
333 global land areas. *Int. J. Climatol.* **37**, 4302–4315. (doi:10.1002/joc.5086)
- 334 35. Upham NS, Esselstyn JA, Jetz W. 2019 Inferring the mammal tree: Species-level sets of


- 335 phylogenies for questions in ecology, evolution, and conservation. *PLoS Biol.* **17**,
336 e3000494. (doi:10.1371/journal.pbio.3000494)
- 337 36. Revell LJ. In press. APPLICATION phytools: an R package for phylogenetic comparative
338 biology (and other things). (doi:10.1111/j.2041-210X.2011.00169.x)
- 339 37. Pagel M. 1999 Inferring the historical patterns of biological evolution. *Nature.* **401**, 877–
340 884. (doi:10.1038/44766)
- 341 38. Genoud M, Isler K, Martin RD. 2018 Comparative analyses of basal rate of metabolism in
342 mammals: data selection does matter. *Biol. Rev.* **93**, 404–438. (doi:10.1111/brv.12350)
- 343 39. Knepper MA, Kwon TH, Nielsen S, Ingelfinger JR. 2015 Molecular physiology of water
344 balance. *N. Engl. J. Med.* **372**, 1349–1358. (doi:10.1056/NEJMra1404726)
- 345 40. Bittner NKJ, Mack KL, Nachman MW. 2020 Plasticity in gene expression facilitates invasion
346 of the desert environment in house mice. *bioRxiv Prepr.* **2011**, 1–9.
347 (doi:<https://doi.org/10.1101/2020.02.10.939231>)
- 348 41. Kim ES, Elbeltagy AR, Aboul-Naga AM, Rischkowsky B, Sayre B, Mwacharo JM, Rothschild
349 MF. 2016 Multiple genomic signatures of selection in goats and sheep indigenous to a hot
350 arid environment. *Heredity (Edinb).* **116**, 255–264. (doi:10.1038/hdy.2015.94)
- 351 42. Wu H *et al.* 2014 Camelid genomes reveal evolution and adaptation to desert
352 environments. *Nat. Commun.* **5**, 10. (doi:10.1038/ncomms6188)
- 353 43. Yang J *et al.* 2016 Whole-Genome Sequencing of Native Sheep Provides Insights into Rapid
354 Adaptations to Extreme Environments. *Mol. Biol. Evol.* **33**, 2576–2592.
355 (doi:10.1093/molbev/msw129)

356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378



379
380 **Figure 1**—Ancestral state reconstruction for mean annual aridity index (AI, left) and maximum urine osmolalities
381 (mOsm/kg; right) in a phylogenetic tree with 121 mammalian species.

382



383

384

385 **Figure 2- Maximum urine osmolality (mOsm/Kg) and mean aridity index (AI).** (A) Boxplot for maximum urine osmolality
386 in endurers, evaders, and evaporators from environments with different degree of aridity, based on 121 different
387 species. Endurers, evaders, and evaporators are classified based on body mass following [5] and aridity classification
388 according to [33]. (B) Scatterplot using mean aridity index as a continuous variable.

389

390

391

392

393

394

395

396

397 **Table 1** -Phylogenetic signal (λ) for variables with significant p-value. Variables for which λ is close to 1, and with p-
398 values < 0.05, are variables with significant phylogenetic signals.

Variable	Page's λ (p-value)
Log10(maximum urine osmolality)	0.988 (3.26e-12)
Log10 (body mass)	1.000(1.26e-41)
Mean annual aridity index	0.854 (0.0002)
Mean annual average temperature	0.668 (0.0026)
Mean temperature of the coldest month	0.772(0.0002)
Mean annual temperature range	0.840(4.90 e-06)

399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430

431 **Table 2 –PGLS models for predicting mammalian log₁₀(maximum urine osmolality).** Model notation refers to the
 432 combination of variables used as predictors while taking phylogeny into account. Highlighted in bold are significant p-
 433 values. Variables are AI = mean aridity index, BM = Log₁₀ body mass (kg), C = condition (dehydrated-D relative to fed-
 434 F, given salt load-S, given protein load-P, unknown-x and combinations of these), M = Method (unknown-? relative to
 435 addition of solutes-A, Freezing point-F, Vapor pressure-V, and combinations of these), Di = Diet (Carnivorous-C relative
 436 to Fructivorous-Fr, Granivorous-G, Herbivorous-H, Insectivorous-I, Omnivorous-O, and combinations of these).
 437

<i>Dataset</i>	<i>Model</i>	<i>AIC</i>	<i>variable</i>	<i>Coefficient</i>	<i>s.e</i>	<i>t-value</i>	<i>p-value</i>
<i>All</i> (<i>n=121</i>)	AI + BM + C + M +Di	-16.3	intercept	3.703	0.097	37.99	0.0000
			AI	-0.368	0.043	-8.47	0.0000
			BM	-0.082	0.015	-5.54	0.0000
			DF _{condition}	0.023	0.080	0.28	0.7763
			DP _{condition}	-0.081	0.108	-0.75	0.4537
			DPS _{condition}	0.020	0.178	0.11	0.9104
			DS _{condition}	-0.047	0.106	-0.45	0.6573
			P _{condition}	-0.006	0.077	-0.08	0.9341
			S _{condition}	0.049	0.065	0.75	0.4536
			SP _{condition}	0.148	0.148	1.00	0.3202
			X _{condition}	-0.049	0.047	-1.06	0.2911
			A _{method}	-0.068	0.153	-0.44	0.6592
			F _{method}	0.016	0.039	0.41	0.6801
			FV _{method}	0.058	0.097	0.60	0.5530
			V _{method}	0.106	0.055	1.93	0.0570
			Cl _{diet}	-0.158	0.141	-1.12	0.2649
			Fr _{diet}	-0.533	0.149	-3.57	0.0005
			G _{diet}	0.013	0.103	0.13	0.8980
			H _{diet}	-0.132	0.088	-1.50	0.1366
I _{diet}	-0.130	0.107	-1.22	0.2260			
O _{diet}	-0.055	0.095	-0.58	0.5634			
<i>Dehydrated</i> (<i>n=87</i>)	AI + BM + M +Di	-34.7	intercept	3.679	0.132	27.83	0.0000
			AI	-0.344	0.043	-7.96	0.0000
			BM	-0.067	0.016	-4.18	0.0001
			A _{method}	-0.107	0.141	-0.75	0.4528
			F _{method}	-0.007	0.042	-0.16	0.8702
			FV _{method}	0.096	0.104	0.93	0.3563
			V _{method}	0.029	0.070	0.42	0.6758
			Cl _{diet}	-0.098	0.165	-0.60	0.5536
			Fr _{diet}	-0.406	0.178	-2.28	0.0254
			G _{diet}	0.090	0.142	0.63	0.5281
			H _{diet}	-0.106	0.135	-0.79	0.4333
			I _{diet}	-0.049	0.140	-0.35	0.7261
			O _{diet}	0.002	0.139	0.02	0.9873

